

Traveler's thrombosis: no fear, just some advice



Garda Aeroporti

Valerio Catullo Airport
Verona, Italy
October 13, 2025

Air travel, especially for long distances, can increase the **risk of venous thromboembolism**. The risk of deep vein thrombosis, mainly related to the prolonged immobility of the lower limbs, is however **limited**. Nevertheless, fear of venous thromboembolism during air travel may lead to improper practices, such as inappropriate self-administration of anticoagulant or antiplatelet drugs. Furthermore, it is forgotten how simple and correct behaviours, such as regular mobilization and adequate hydration, can reduce the risk of venous thromboembolism and allow to travel safely.



WORLD THROMBOSIS DAY
13 OCTOBER

We invite you to learn more about “**traveler's thrombosis**” so as **not to be afraid of it** and **prevent it with some straightforward precautions**

Traveler's thrombosis: no fear, just some advice

Venous thrombosis is due to the activation of coagulation in the wrong place, inside the venous vessels, causing their **obstruction by a blood clot**. Venous thromboembolism includes **deep vein thrombosis**, generally in the lower limbs, and **pulmonary embolism**, which is usually caused by a piece of clot, the embolus, which detaches from the venous thrombus, reaches the heart and then the lungs, obstructing the flow of blood to the lungs. Venous thromboembolism can be triggered by **multiple risk factors including long-distance air travel**.



Causes

The causes of traveler's thrombosis are diverse and often unclear. Although **various mechanisms** have been proposed, from dehydration causing blood thickening to exposure to reduced oxygen pressure during air travel, the **primary risk factor is certainly prolonged immobility** (which must be taken into account not only during air travel, but also while on other forms of travelling such as car or train journeys). **Prolonged sitting**, especially in **confined spaces**, limits leg movement and reduces blood flow in the veins of the legs, thus promoting the risk of thrombosis.

How it presents

Symptoms of deep vein thrombosis: pain, redness and/or swelling of the lower limbs (much more rarely the upper limbs), usually occurring asymmetrically.

Symptoms of pulmonary embolism: difficulty breathing and shortness of breath (the most important symptom), chest pain, heart palpitations, fainting, coughing up traces of blood.

The size of the problem

The absolute **risk** of venous thrombosis in the air traveler is **generally low** but **increases with the duration of the journey**. It has been documented when the journey lasts more than 4 hours and is greater for journeys longer than 10 hours. Studies have estimated that air travel lasting more than 4 hours is associated with a 2-3 times higher risk of venous thrombosis compared to the general population, nevertheless leading to only a moderate probability (with approximately 1 thrombotic event occurring per 4,600 flights, or 0.02%).

Who is most at risk

The risk of traveler's thrombosis is higher in individuals who already have other risk factors:

- age > 40 years
- obesity
- female gender
- pregnancy and puerperium (6-8 weeks postpartum)
- use of oral contraceptives and hormone replacement therapies
- personal history of previous venous thromboembolism
- family history of venous thromboembolism (particularly in the case of first-degree family members who experienced the thrombotic event at a young age)
- hereditary thrombophilias
- active cancer
- smoking
- recent surgery or trauma (within 6 weeks prior to travel)
- other specific medical conditions (e.g. chronic heart failure, rheumatological diseases, venous insufficiency in the lower limbs).



What I can do to reduce thrombotic risk

1. While waiting for boarding, **walk** through the permitted areas.
2. During the journey, **if possible, get up every two hours and walk down the aisle**. When sitting, alternately flex and extend your knees and ankles at regular intervals and move your shoulders.
3. On board, **sit stretching your legs** to avoid keeping your knees flexed at 90 degrees (this position can hinder blood flow in the veins of your legs).
4. Minimize the size of carry-on luggage under the seat to ensure that your legs have **enough room to move**.
5. On the day of the journey, wear **comfortable clothes**.
6. Before and during the journey, **keep well hydrated** with water and **abstain from alcohol**.
7. **Avoid taking sleeping pills**, which can promote long periods of immobility.

Special situations

1. In case of **reduced mobility and/or high risk** of venous thromboembolism, consider the use of **graduated compression stockings** to promote blood flow from the veins of the legs to the heart, if not contraindicated and after prior medical consultation.
2. **Very rarely**, for some travelers at **high thrombotic risk**, **anticoagulant or antiplatelet drugs** to be administered/taken before travel may be considered. The use of these drugs remains controversial and with limited evidence for the prevention of traveler's thrombosis. Therefore, **their self-administration is strongly discouraged in the absence of specific medical evaluation and advice**, also in consideration of the possible side effects (e.g. bleeding and allergies).
3. The traveler may **already be on anticoagulant or antiplatelet therapy** for specific medical conditions with particularly high thrombotic risk. **The previous recommendations** (see the paragraph “What I can do to reduce the thrombotic risk”) **remain valid also in this case**.



Scan the QR code!



UNIVERSITÀ di VERONA

Dipartimento di MEDICINA



AZIENDA OSPEDALIERA UNIVERSITARIA INTEGRATA VERONA



Verona Aeroporti

Garda Aeroporti

Supported by an unrestricted educational sponsorship



Thrombosis &
Coagulation Group
Internal Medicine B
Verona, Italy